

Example MEDICATION LIST

Discontinue the use of any medications that may affect the outcome of allergy testing as described on the previous page. Drugs which may affect skin testing, include ANTIHISTAMINES, (including allergy, cold & sinus medicine, both prescription and non-prescription), and certain ANTIDEPRESSANTS.

The ANTIHISTAMINES listed below should be discontinued for **4 DAYS** before skin testing. The ANTIDEPRESSANTS listed should be discontinued for **1 DAY** before skin testing. (If you have any questions regarding medicines you are taking that are not listed, please call the office):

Allegra(Fexofenadine)	Contac	Rescon
Allegra-D	Coricidin	Rescon Jr.
Alavert	DA Chewable	Robitussin Night Time Cold
Alavert-D	DA II	Ryna
Actifed	Deconamine	Rynatan
Alka-Seltzer Plus/Cold	Dimetapp	Rynatuss
Allerest	Dramamine	Semprex-D(Acrivastine)
Antivert(Meclizine)	Drixoral	Sine-Off Night Time
BC Allergy Sinus	Dura-Vent DA	Sinutab
Bonine	Excedrin PM	Sudafed Cold & Allergy
Bromfed	Extendryl	Tavist(Clemastine)
Children's Tylenol Allergy-D	Fedahist	Tavist-D
Children's Tylenol Multisymptom Cold, Chewable & Liquid	Kronofed	Thera-Flu
Chlortrimeton	Librium	Triaminic
Chlorpheniramine Maleate	Nolahist	Tussi-12
Clarinox(Desloratadine)	Nyquil	Tussionex
Clarinox D	PBZ-SR	Tylenol Allergy Sinus
Claritin(Loratadine)	PediaCare	Tylenol Flu Night Time
Claritin-D	Periactin	Tylenol PM
Comtrex	Phenergan	Vicks 44M
	Pyribenzamine	

IF POSSIBLE, PLEASE DISCONTINUE ANY ANTI-DEPRESSANTS **24 HOURS** prior.

ASTELIN, ASTEPRO, AZELASTINE, PATANASE AND DYMISTA NASAL SPRAYS and BENADRYL LIQUID & TABLETS (DIPHENHYDRAMINE) must be discontinued for **2 DAYS**.

ATARAX, HYDROXYZINE, VISTARIL, ZYRTEC, ZYRTEC-D, CETIRIZINE, XYZAL, LEVOCETIRIZINE, DOXEPIN AND SINEQUAN should be discontinued for **7 DAYS**.

There is no need to discontinue Corticosteroid Nose Sprays, (Flonase, Nasacort, Nasarel, Nasonex, Rhinocort, Veramyst, Fluticasone, etc.).